

1218 -MEAT EXTRACT (STD) TBL POWDER (Culture Media Ingredient)

INTENDED USE

A nutritious extract used as an ingredient in the preparation of culture media for cultivation of a wide variety of fastidious microorganisms.

PRODUCT SUMMARY AND EXPLANATION

Meat Extract (STD) TBL Powder is derived from infusion of meat and provides an undefined source of nutrients. It is not exposed to the harsh treatment used for protein hydrolysis, so it can provide some of the nutrients lost during peptone and amino acids, nucleotide fractions, organic acids, minerals and some vitamins. Manufacture. Meat Extract (STD) TBL Powder is a mixture of peptides

PRINCIPLE

Meat Extract (STD) TBL Powder is manufactured from meat with low fat content and can be considered as complementing the nutritive properties of peptone by contributing minerals, phosphates, energy sources and those essential factors missing from peptone. Meat Extract (STD) TBL Powder powder is a source of amino acids, peptides and other nutrients.

INSTRUCTION FOR USE

Meat Extract (STD) TBL Powder is frequently used at a concentration of 0.3 to 1.0% in culture media, although concentrations may vary depending on the nutritional requirements for the medium formulation.

QUALITY CONTROL SPECIFICATIONS

Appearance	:	Creamish to white colour free flowing powder having characteristic odour but not pungent smell.
Solubility (2% soln. at 25°C)	:	Completely soluble in distilled water, clear.
Clarity (2% Soln. at 121°C)	:	Absolute clear solution. No ppt..
pH (2% Soln. at 25°C)	:	6.5 – 7.5
Loss on drying (at 105°C)	:	NMT – 6.0%
Total Nitrogen (DWB)	:	NLT – 12.5%
α-Amino Nitrogen	:	NLT – 3.5%
Total Ash	:	NMT – 15.0%
Chloride (as NaCl)	:	NMT – 6.0%
Microbial parameters	:	Passes Test
Indole Test	:	Positive

INTERPRETATION

Cultural Characteristic observed in 2% Meat Extract (STD) TBL Powder and 1.5% agar after incubation at 35-37°C for 18-48 hours.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth
<i>Staphylococcus aureus</i>	25923	50-100	Good - Luxuriant
<i>Escherichia coli</i>	25922	50-100	Good - Luxuriant
<i>Pseudomonas aeruginosa</i>	27853	50-100	Good - Luxuriant
<i>Clostridium perfringens</i>	12924	50-100	Good - Luxuriant
<i>Enterococcus faecalis</i>	29212	50-100	Good - Luxuriant
<i>Streptococcus pyogenes</i>	19615	50-100	Good - Luxuriant

PACKAGING:

Standard packing is 500gm in plastic bottle. After packing tightly closed in a dry and well-ventilated place.

STORAGE

Store at room temperature in cool place, Keep plastic bottle tightly closed in a dry and well-ventilated place. Use before expiry date on label. On opening, product should be properly stored in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use.


Product Deterioration: Do not use product if any contamination, discoloration or other sign of deterioration is found.

DISPOSAL

After use, contact a licensed professional waste disposal service to dispose of this material. Dispose of as unused product.

REFERENCES

1. United States Pharmacopeial Convention. 2006. The United States pharmacopeia 29/The national formulary 24—2006. United States Pharmacopeial Convention, Inc., Rockville, Md.
2. Cote. 1999. Media composition, microbial, laboratory scale. In Flickinger and Drew (ed.), Encyclopedia of bioprocess technology: fermentation, biocatalysis, and bioseparation. John Wiley & Sons, Inc., New York.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 05th Oct. 2019