

1509V – VEG. PEPTONE SPECIAL (Culture Media Ingredient)

INTENDED USE

Veg.Peptone Special used in the preparation of culture media employed for cultivation of a wide variety of microorganisms.

PRODUCT SUMMARY AND EXPLANATION

Veg.Peptone Special is used in preparing microbiological culture media and in producing bacterial toxins and also usable in synthetic media in acclimatization of microorganisms in bioreactor studies. It's support to growth of Staphylococci, Streptococci, Pneumococci and also suitable for isolating and cultivating Haemophilus and Neisseria. It is off white to Creamish yellow colour, free flowing powder having characteristic odour but not pungent smell. It is completely soluble in distilled Water, Clear. Insoluble in alcohol.

PRINCIPLE

Veg.Peptone Special is enzymatic digest of protein used in preparing microbiological culture media and in producing bacterial toxins. Proteose peptone provide nitrogen in a form that is readily available for bacterial growth. It is superior in nutritious of fastidious microorganism.

INSTRUCTION FOR USE

Veg.Peptone Special is used in media for the production of bacterial toxins. It is used in preparing chocolate agar for propagating of Neisseria species. It is also used for the cultivation of bacteria with high nutritional requirements, as for example Haemophilus, Salmonella, staphylococcus etc. species.

QUALITY CONTROL SPECIFICATIONS

Light yellowish to brownish yellow colour, free flowing powder **Appearance**

having characteristic odour but not pungent smell.

Completely soluble in distilled Water, Clear. Insoluble in alcohol. Solubility (2% Soln. at 25°C) :

6.5 - 7.5pH (2% Soln. at 25 °C) Loss on drying (at 105 °C) NMT - 5.0% NLT - 12.5% Total Nitrogen (DWB) α-Amino Nitrogen NLT - 3.5%**Total Ash** NMT - 15.0% Chloride (as NaCl) NMT - 5.0% **Indole Test** Positive **Microbial Parameter Passes Test Growth Promotion Test Passes Test**

INTERPRETATION

Cultural Characteristic observed in 2% Veg. PP Peptone and 1.5% agar after incubation at 35-37°C for 18-24 hours.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth
Staphylococcus aureus	6538	50-100	Luxuriant
Escherichia coli	8739	50-100	Luxuriant
Pseudomonas aeruginosa	9027	50-100	Luxuriant
Enterobacter aerogenes	13048	50-100	Luxuriant
Salmonella typhi	6539	50-100	Luxuriant
Streptomyces albus	3004	50-100	Luxuriant











Streptococcus pyogenes	19615	50-100	luxuriant w/ beta haemolysis (With addition of sterile 5% sheep blood to above medium, after an incubation at 35-37°C for 48 hours
Neisseria gonorrhoeae	19424	50-100	luxuriant w/ beta haemolysis (With addition of sterile 10% sheep blood to above medium heated to 80-90°C until blood has turned to chocolate brown and incubated in 10% CO2 atmosphere at 35-37°C for 48 hours).

PACKAGING:

Standard packing is 500gm in plastic bottle. After packing tightly closed in a dry and well-ventilated place.

STORAGE

Keep plastic bottle tightly closed in a dry and well-ventilated place, Store in cool place. Use before expiry date on label. On opening, product should be properly stored in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the plastic bottle after use.


Product Deterioration: Do not use product if any contamination, discoloration or other sign of deterioration is found.

DISPOSAL

After use, contact a licensed professional waste disposal service to dispose of this material. Dispose of as unused product.

REFERENCES

- 1.Kirkbride, Berthelsen and Clark. 1931. Comparative studies of infusion and infusion-free diphtheria toxin in antitoxin production and in standardization by the flocculation, subcutaneous, and intracutaneous tests. J. Immunol. 21:1-20.
- 2. Hazen and Heller. 1931. Further studies upon the effect of various carbohydrates on production of diphtheria toxin with special reference to its flocculating titer and final pH. J. Bacteriol. 23:195-209.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only

Revision: 05th Oct. 2019









