

TM 1080 - SABOURAUD DEXTROSE MALTOSE BROTH

INTENDED USE

For cultivation of molds, yeasts & aciduric organisms as well as testing antimycotic substances.

PRODUCT SUMMARY AND EXPLANATION

Sabouraud Dextrose Agar is Carliers modifications of the formulation described by Sabouraud for the cultivation of fungi, particularly those associated with skin infections. Sabouraud Dextrose Maltose Broth is used for the cultivation of yeast, moulds and other aciduric organisms.

Sabouraud dextrose media are peptone media supplemented with dextrose to support the growth of fungi. The acid reaction of the final medium is inhibitory to a large number of bacteria making it particularly useful for cultivating fungi and aciduric microorganisms. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously. Incubate cultures for 4 to 6 weeks before reporting as negative.

COMPOSITION

Ingredients	Gms / Ltr		
Casein enzymic hydrolysate	5.000		
Peptic digest of animal tissue	5.000		
Dextrose	10.000		
Maltose	10.000		

PRINCIPLE

Casein enzymic hydrolysate and peptic digest of animal tissue provide nitrogen, vitamins, minerals, amino acids and growth factors. Dextrose and maltose provide an energy source for the growth of microorganisms. The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens.

INSTRUCTION FOR USE

- Dissolve 30.0 grams in 1000 ml distilled water.
- Heat if necessary to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes, do not overheat.
- Mix well and dispense as desired.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder : Cream to yellow homogeneous free flowing powder.

Appearance of prepared medium : Light amber coloured clear solution in tubes.

pH (at 25°C) : 5.4±0.2

INTERPRETATION

Cultural characteristics observed after an incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Incubation Temperature	Incubation Period	









Aspergillus brasiliensis	16404	10-100	Luxuriant	25-30°C	Upto 5 days
Candida albicans	10231	10-100	Luxuriant	25-30°C	Upto 5 days
Pencillium notatum	10108	10-100	Luxuriant	25-30°C	Upto 5 days
Trichophyton mentagrophyts	9533	10-100	Luxuriant	25-30°C	Upto 5 days
Trichophyton gallinae	22243	10-100	Luxuriant	25-30°C	Upto 5 days
Trichophyton rubrum	28191	10-100	Luxuriant	25-30°C	Upto 5 days
Trichopyton ajelloi	24885	10-100	Luxuriant	25-30°C	Upto 5 days
Escherichia coli	25922	50-100	Good- luxuriant(inhibit ed on media with low pH)	25-30°C	Upto 5 days
Lactobacillus casei	9595	50-100	Good- luxuriant	25-30°C	Upto 5 days
Saccharomyces cerevisiae	9763	10-100	Good- luxuriant	25-30°C	Upto 5 days

PACKAGING:

In pack size of 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.













Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- 1. Carlier G. I. M., 1984, Brit. J. Derm. Syph., 60:61
- 2. Sabouraud R., 1892, Ann. Dermatol. Syphil. 3: 1061.
- 3. Merkblatt 18: Verpackgs- Rdsch, 1974, 25/1: Techn- Wiss. Beilage, 5-8
- 4. Merkblatt 19: Verpackgs- Rdsch, 1974, 25/6: Techn- Wiss. Beilage, 569-575
- 5. Merkblatt 21: Verpackgs- Rdsch, 1974, 25/7: Techn- Wiss. Beilage, 53-55
- 6. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Yolken R. H., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only
Revision: 08 Nov., 2019







