

TM 215 - MIDDLEBROOK 7H11 AGAR BASE W/O MALACHITE GREEN

INTENDED USE

For isolation, cultivation and determination of antimicrobial susceptibility of Mycobacteria.

PRODUCT SUMMARY AND EXPLANATION

Middlebrook 7H11 Agar Base w/o Malachite Green is based on Middlebrook 7H11 Agar which is a modification of Middlebrook 7H10 Agar used for the isolation, cultivation and sensitivity testing of Mycobacterium tuberculosis. It was shown by Cohn et al that the addition of casein enzymic hydrolysate enhanced the growth of more fastidious Mycobacterium tuberculosis strains which in turn was helpful in drug susceptibility testing.

COMPOSITION

Ingredients	Gms / Ltr	
Casein enzymic hydrolysate	1.000	
Ammonium sulphate	0.500	
Monopotassium phosphate	1.500	
Disodium phosphate	1.500	
Sodium citrate	0.400	
Magnesium sulphate	0.050	
L-Glutamic acid	0.500	
Ferric ammonium citrate	0.040	
Pyridoxine	0.001	
Biotin	0.0005	
Agar	15.000	

PRINCIPLE

The media consists of many inorganic salts which help for growing Mycobacteria. Citric acid formed from sodium citrate helps in retaining inorganic cations in solution. Glycerol supplies carbon and energy. OADC Supplement contains oleic acid, bovine albumin, sodium chloride, dextrose and catalase. Oleic acid and other long chain fatty acids are metabolized by Mycobacteria. Dextrose is an energy source. Catalase neutralizes toxic peroxides, while albumin protects tubercle bacilli from toxic agents. Malachite green partially inhibits other bacteria.

INSTRUCTION FOR USE

- Dissolve 10.25 grams in 450 ml distilled water containing 2.5 ml glycerol.
- Heat to boiling to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Cool to 50°C. Aseptically add contents of 1 vial of Middlebrook OADC Growth Supplement.
- Mix thoroughly before dispensing.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder : Light yellow to light green homogeneous free flowing powder.

Appearance of prepared medium : Light amber coloured clear to slightly opalescent gel with greenish tinge.

pH (at 25°C) : 6.6±0.2









INTERPRETATION

Cultural characteristics observed after an incubation.

Microorganism	АТСС	Inoculum (CFU/ml)	Growth	Recovery	Incubation Temperature	Incubation Period
Mycobacterium tuberculosis	25618	50-100	Good- luxuriant	>=50 %	35-37°C	2-4 weeks
Mycobacterium fortuitum	6841	50-100	Good- luxuriant	>=50 %	35-37°C	2-4 weeks
Mycobacterium smegmatis	14468	50-100	Good- luxuriant	>=50 %	35-37°C	2-4 weeks

PACKAGING:

In pack size of 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.


Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- 1. Middlebrook and Cohn, 1958, Am. J. Public Health, 48:844.
- 2. 2.Cohn et.al, 1968, Am.Rev.Resp.Dis., 98:295.
- $3.\ 3. Mac Faddin\ J.F.,\ 1985,\ Media\ for\ Isolation-Cultivation-Identification-Maintenance\ of\ Medical\ Bacteria,\ Vol.\ I,\ Williams\ and\ Wilkins,\ Baltimore$



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.















*For Lab Use Only Revision: 08 Nov., 2019









