

## TM 2174 – LYSINE INDOLE MOTILITY MEDIUM, MODIFIED

### INTENDED USE

Is used as an aid for the identification of members of *Enterobacteriaceae* on the basis of lysine decarboxylase, indole production and motility.

### PRODUCT SUMMARY AND EXPLANATION

Lysine Indole Motility Medium, Modified is a semisolid medium used for the differentiation of *Enterobacteriaceae* group by lysine decarboxylation, indole production and motility.

### COMPOSITION

Ingredients	Gms / Ltr
Peptone	12.800
Yeast extract	3.000
L-Tryptophan	0.500
L-Lysine	10.000
Dextrose (Glucose)	1.000
Bromocresol purple	0.020
Agar	2.700

### PRINCIPLE

This medium consists of Peptone and yeast extract which supply amino acids and other complex nitrogenous substances. Dextrose (Glucose) is a source of energy. A small amount of agar is added for demonstration of motility along the stab line of inoculation. Growth of motile organisms extends out from the line of inoculation, while non-motile organisms grow only along the stab line. Bromocresol purple serves as the pH indicator.

When inoculated with an organism that ferments dextrose, acids are produced that lower the pH, causing the indicator in the medium to change from purple to yellow. The acidic pH also stimulates decarboxylase enzyme activity. Organisms that possess a specific decarboxylase degrade the amino acid provided in the medium, yielding a corresponding amine. Lysine decarboxylation yields cadaverine. The production of these amines elevates the pH and causes the medium in the bottom portion of the tube to revert to a purple color. The medium in the upper portion of the tube remains acidic because of the higher oxygen tension. If the organism being tested does not produce the required decarboxylase, the medium remains yellow (acidic) throughout or yellow with a purple or red reaction near the top. Indole is produced in this medium by organisms that possess the enzyme tryptophanase. Tryptophanase degrades tryptophan, yielding indole. It can be detected in the medium by adding Kovacs reagent to the agar surface. Indole combines with the p-dimethylaminobenzaldehyde of Kovacs reagent and produces a red complex.

### INSTRUCTION FOR USE

- Dissolve 30.02 grams in 1000 ml purified/distilled water.
- Heat to boiling to dissolve the medium completely.
- Dispense into tubes in 5 ml amounts and sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Cool the tubes in an upright position.

### QUALITY CONTROL SPECIFICATIONS



**Appearance of Powder** : Cream to greenish yellow homogeneous free flowing powder.  
**Appearance of prepared medium** : Reddish purple coloured clear to slightly opalescent gel forms in tubes as butts.  
**pH (at 25°C)** : 6.7 ± 0.2

## INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Motility	Indole production	Lysine decarboxylase	Incubation Temperature	Incubation Period
<i>Klebsiella aerogenes</i>	13048	50-100	Positive, growth away from stabline	Negative reaction	Positive reaction, purple colour	35-37°C	18-24 Hours
<i>Escherichia coli</i>	25922	50-100	Positive, growth away from stabline	Positive, red ring at interface of the medium on addition of Kovac's reagent	Positive reaction, purple colour	35-37°C	18-24 Hours
<i>Klebsiella pneumoniae</i>	13883	50-100	Negative, growth along the stabline	Occasional reaction	Positive reaction, purple colour	35-37°C	18-24 Hours
<i>Proteus mirabilis</i>	25933	50-100	Positive, growth away from stabline	Negative reaction	Negative reaction	35-37°C	18-24 Hours
<i>Proteus vulgaris</i>	13315	50-100	Positive, growth away from stabline	Positive, red ring at interface of the medium on addition of Kovac's reagent	Negative reaction	35-37°C	18-24 Hours
<i>Salmonella Enteritidis</i>	13076	50-100	Positive, growth away from stabline	Negative reaction	Positive reaction, purple colour	35-37°C	18-24 Hours
<i>Shigella flexneri</i>	12022	50-100	Negative, growth along the stabline	occasional reaction	Negative reaction	35-37°C	18-24 Hours

## PACKAGING:

In pack size of 500 gm bottles.

## STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

**Product Deterioration:** Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.















## DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

## REFERENCES

1. Igarashi.H et al.(1969)A new differential medium for enteric pathogens, lysine-indole-motility medium
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

 <b>GMP</b> Good Manufacturing Practices Certified	 <b>IVD</b> For In Vitro Diagnostic Use	 <b>QTY.</b> Quantity	 <b>LOT/ B. NO.</b> Lot / Batch Number	 <b>REF</b> Catalogue Number	 <b>Manufacturer</b>
 <b>Temperature Unit</b>	 <b>EC REP</b> Authorized Representative <small>MedNet GmbH Birkstrasse 10, 48163 Münster, Germany</small>	 <b>European Conformity</b>	 <b>QR Code</b>	 <b>Consults Instructions for Use</b>	 <b>Best Before</b>

**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

**\*For Lab Use Only**  
**Revision: 08 Nov., 2019**