

TM 2320 - SABOURAUD DEXTROSE AGAR MEDIUM W/ ANTIBIOTICS (as per IP)

INTENDED USE

For selective cultivation of yeasts and moulds.

PRODUCT SUMMARY AND EXPLANATION

Sabouraud Dextrose Agar Medium w/antibiotics is recommended for cultivation of yeasts and moulds by Indian Pharmacopoeia. This medium was described originally by Sabouraud for the cultivation of fungi, particularly useful for the fungi associated with skin infections. The medium is used with antibiotics such as tetracycline and benzylpenicillin for the isolation of pathogenic fungi from materials containing large numbers of fungi or bacteria. Some pathogenic fungi may produce infective spores which are easily dispersed in air, so examination should be carried out in safety cabinet.

COMPOSITION

Ingredients	Gms / Ltr
Peptones (meat and casein)	10.000
Dextrose monohydrate	40.000
Agar	15.000

PRINCIPLE

Peptic digest of animal tissue and pancreatic digest of casein provide nitrogenous compounds. Dextrose provides an energy source. Tetracycline and benzyl penicillin inhibits a wide range of Gram-positive and Gram-negative bacteria which makes the medium selective for fungi. The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens.

INSTRUCTION FOR USE

- Dissolve 65 grams in 1000 ml distilled water.
- Heat to boiling to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Aseptically add rehydrated contents of one vial of Tetracycline Selective Supplement. Mix well before pouring into sterile Petri plates.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder	: Cream to yellow homogeneous free flowing powder.
Appearance of prepared medium	: Light amber coloured clear to slightly opalescent gel forms in Petri plates.
pH (at 25°C)	: 5.40-5.80

INTERPRETATION

Cultural response was carried out in accordance with IP, with added Tetracycline Selective Supplement, after an incubation.



Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Incubation Temperature	Incubation Period
<i>Candida albicans</i>	10231	10 -100	Luxuriant (white colonies)	≥ 70 %	20-25°C	≤ 5 days
<i>Aspergillus brasiliensis</i>	16404	10 -100	Luxuriant	≥ 70 %	20-25°C	≤ 5 days
<i>Candida albicans</i>	2091	10 -100	Luxuriant	≥ 70 %	20-25°C	≤ 5 days
<i>Saccharomyces cerevisiae</i>	9763	10 -100	Luxuriant	≥ 70 %	20-25°C	≤ 5 days
<i>Escherichia coli</i>	25922	$\geq 10^3$	Inhibited	0 %	20-25°C	≤ 5 days
<i>Escherichia coli</i>	8739	$\geq 10^3$	Inhibited	0 %	20-25°C	≤ 5 days
<i>Escherichia coli</i>	9002	$\geq 10^3$	Inhibited	0 %	20-25°C	≤ 5 days
<i>Trichophyton rubrum</i>	28191	10 -100	Good	40-50%	20-25°C	≤ 5 days
<i>Lactobacillus casei</i>	334	$\geq 10^3$	Inhibited	0 %	20-25°C	≤ 5 days

PACKAGING:

In pack size of 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.




Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

1. Indian Pharmacopoeia 2010, Ministry of Health and Family welfare, Government of India, New Delhi
2. Sabouraud K., 1892, Ann. Dermatol. Syphilol, 3:1061.
3. Ajello L., 1957, J. Chron. Dis., 5:545.
4. Lorian (Ed.), 1980, Antibiotics in Laboratory Medicine, Williams and Wilkins, Baltimore.
5. Murray, P. R 2005, In Manual of Clinical Microbiology, 7th ed., ASM, Washington, D.C.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 08 Nov., 2019