

TMV 330 – PEPTONE WATER (VEG.)

INTENDED USE

General purpose growth medium and as the base of carbohydrate fermentation media.

PRODUCT SUMMARY AND EXPLANATION

Peptone Water (Veg) is prepared by replacing peptic digest of animal tissue with Veg peptone which is free from BSE/TSE risks. Peptone Water (Veg) is the modification of Peptone Water which is recommended for studying the ability of an organism to ferment a specific carbohydrate and indole production, which aid in differentiation of genera and species. Peptone Water (Veg) with pH adjusted to 8.4 is suitable for the cultivation and enrichment of *Vibrio* species.

COMPOSITION

Ingredients	Gms / Ltr
Veg Peptone	10.000
Sodium chloride	5.000

PRINCIPLE

The medium consists of Veg peptone which provides nitrogenous and carbonaceous compounds, long chain amino acids, vitamins provides essential nutrients. Sodium chloride maintains the osmotic balance of the medium.

INSTRUCTION FOR USE

- Dissolve 15.0 grams in 1000 ml distilled water.
- Heat if necessary to dissolve completely.
- Dispense in tubes with or without inverted Durhams tubes and sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder	: Light yellow coloured, may have slightly greenish tinge homogeneous, free flowing powder.
Appearance of prepared medium	: Light yellow coloured, clear solution without any precipitate.
pH (at 25°C)	: 7.2 ± 0.2

INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Indole test	Acid production	Incubation Temperature	Incubation Period
<i>Staphylococcus aureus</i>	25923	50-100	Good-Luxuriant	-	Acid production, yellow colouration of the medium	35-37°C	18-24 Hours



<i>Escherichia coli</i>	25922	50-100	Good-Luxuriant	Positive reaction, red ring observed on addition of Kovac's Indole reagent	Acid production, yellow colouration of the medium	35-37°C	18-24 Hours
<i>Salmonella serotype Typhimurium</i>	14028	50-100	Good-Luxuriant	-	Acid production, yellow colouration of the medium	35-37°C	18-24 Hours

PACKAGING:

In pack size of 100 gm and 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.













Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

1. MacFaddin J.F., 2000(ed), Biochemical Tests for Identification of Medical Bacteria, 3rd edition, Lippincott Williams and Wilkins, New York
2. Finegold and Baron, 1986, Bailey and Scotts Diagnostic Microbiology, 7th ed., The C.V. Mosby Co., St. Louis.
3. Patrick R. Murray, Baron, Pfaller, Tenover and Tenover (Eds.), 2005, In Manual of Clinical Microbiology, 7th ed., ASM, Washington, D.C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedNet GmbH Borkstrasse 10, 48163 Moenster, Germany</small>	 CE European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 08 Nov., 2019