

TMV 733 – FLUID CASEIN DIGEST SOYA LECITHIN MEDIUM (DOUBLE PACK) (VEG.)

INTENDED USE

For sanitary testing of surfaces.

PRODUCT SUMMARY AND EXPLANATION

This medium is prepared by using Veg hydrolysate is place of Casein enzymic hydrolysate which makes the medium free of BSE/TSE risks. Fluid Casein Digest Soya Lecithin Veg Medium is recommended for sanitary examination of surfaces. Weber and Black had described the importance of a highly nutritional medium containing the neutralizing agents for quaternary ammonium compounds.

COMPOSITION

Ingredients	Gms / Ltr				
Part I					
Veg hydrolysate	20.000				
Soya lecithin	5.000				
Part II					
Polysorbate 20	40.000				

PRINCIPLE

The medium consists of Veg hydrolysate, which provides necessary nutrients for the growth of the organisms. Soya lecithin neutralizes the quaternary ammonium compounds while polysorbate 20 neutralizes phenolic disinfectants, hexachlorophene and formalin.

INSTRUCTION FOR USE

- Dissolve 25.0 grams of Part I in 960 ml purified / distilled water.
- Heat if necessary to dissolve the medium completely.
- Add 40 ml of Part II. Mix well and sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder : Part I : Yellow coloured, may have greenish tinge, homogeneous, free flowing

powder. Part II: Colourless, clear, viscous liquid.

Appearance of prepared medium : Yellow coloured, clear solution without any precipitate.

pH (at 25°C) : 7.3 ± 0.2

INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism ATCC Inoculum Growth Incubation Incubation Temperature Period









Candida albicans	10231	10-100	Good-luxuriant	25-30°C	24-48 Hours
Bacillus subtilis	6633	50-100	Good-luxuriant	35-37°C	18-24 Hours
Escherichia coli	25922	50-100	Good-luxuriant	35-37°C	18-24 Hours
Staphylococcus aureus subsp. aureus	25923	50-100	Good-luxuriant	35-37°C	18-24 Hours

PACKAGING:

In pack size of 100 gm and 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.


Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- 1. Weber and Black, 1948, Soap and Sanitary Chemicals, 24:134.
- 2. Weber and Black, 1948, Am. J. Public Health, 38:1405.
- 3. Favero (chem.), 1967, Microbiological Sampling of Surfaces, Biological Contamination Control Committee, American Asso. for Contamination Control.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only
Revision: 08 Nov., 2019





