

TP 085 – GAMBORG B5 MEDIUM (W/CaCl₂, Vitamins, Agar and Sucrose, W/O IAA & Kinetin)

INTENDED USE

Supports or facilitate plant growth and/or shoot proliferation in two or more plant tissue cultures (both in monocotyledons and dicotyledons).

COMPOSITION

Ingredients	Mg / Ltr
Copper sulphate.5H ₂ O	0.025
Cobalt chloride.6H ₂ O	0.025
Ferric sulphate. 7H ₂ O	27.80
EDTA disodium salt.2H ₂ O	37.30
Boric acid	3.000
Potassium iodide	0.750
Molybdic acid (sodium salt).2H ₂ O	0.25
Calcium chloride	150.00
Manganese sulphate	10.000
Zinc sulphate. 7H ₂ O	2.00
Sodium hydrogen phosphate	130.440
Potassium nitrate	2500.000
Magnesium sulphate	122.09
Ammonium sulphate	134.000
Sodium phosphate monobasic	130.42
Myo-Inositol	100.000
Thiamine HCl	10.00
Nicotinic acid	1.000
Pyridoxine HCl	1.000
Sucrose	20000.00
Agar	8000.00

Formula weight: 31.23 gms/ltr

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder	: White to off white with homogenous mixture of free flowing powder.
Appearance of prepared medium	: Colorless to slight yellow solution, clear, complete.
pH (at 25°C)	: 5.8 ± 0.2

INSTRUCTION FOR USE

Dissolve 31.23 Gms of dehydrated medium in 600 ml of distilled or deionized water at room temperature (15 - 30°C). Rinse media vial with small quantity of distilled water to remove traces of powder. Add the desired heat stable












supplements prior to autoclaving. Continue stirring until the powder has dissolved. Sometimes media does not dissolve completely unless the pH is reduced. For these, lower the pH to about 3.0 to facilitate dissolution of media. The pH of medium is adjusted by using 1N HCl/ 1N NaOH/ 1N KOH. Make up the final volume to 1000ml with distilled water. Mix gently, heat and rotate between intervals until the solution becomes clear. Do not boil, reheat and allow cooling below 50°C during dispensing. Dispense the medium into suitable containers, plug or cap, then autoclave at 15psi (121°C) for 15 minutes, using a slow exhaust cycle. Higher temperatures and/or longer times are not recommended. Cool the autoclaved culture vessels containing medium to 45-50°C and aseptically add desired sterile heat-labile substrates.

Note: Media should be prepared according to formula mentioned on the label however, it is recommended to use an entire container at once. Heat-labile substrates should be added, after autoclaving.

STORAGE

Dehydrated plant tissue culture media is hygroscopic and should be protected from sunlight and moisture. Store the prepared medium at 2–8°C away from direct light. Medium should be used before the expiry date. The entire volume of each bottle should be used immediately after opening or else the unused portion should be stored in desiccators and refrigerated at 2–8°C.

 Good Manufacturing Practices Certified	 Best Before	 Quantity	 Catalogue Number	 Manufacturer
 Temperature Unit	 Lot / Batch Number	 Consults Instructions for Use	 QR Code	

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 5 March., 2022