

TS 003 – POTASSIUM TELLURITE 3.5% (1 ml/vl)

INTENDED USE

For selective isolation of Staphylococci and *Corynebacteria*.

COMPOSITION

Ingredients	Concentration
Potassium tellurite	0.350g
Distilled water	1ml

(per vial sufficient for 200 ml medium)

Applied in Medium

TM 358 - BAIRD PARKER AGAR BASE

TMV 358 - BAIRD PARKER AGAR BASE(VEG.)

TM 635 - BAIRD PARKER AGAR BASE (IS: 5887 (Part II) 1976, reaffirmed 2005.)

TM 943 - BAIRD PARKER AGAR BASE W/ SULPHA

TM 118 - GIOLITTI-CANTONI BROTH BASE

TM 118A - GIOLITTI-CANTONI BROTH BASE (ISO 6888-3:2003)


TM 522 - HOYLE MEDIUM BASE

INSTRUCTION FOR USE

Warm up the refrigerated contents of one vial to 45-50°C. Add aseptically 9 ml sterile distilled water, mix well and add 3 ml in 950 ml sterile, molten, cooled (45-50°C) TM 358 - BAIRD PARKER AGAR BASE / TMV 358 - BAIRD PARKER AGAR BASE(VEG.) / TM 635 - BAIRD PARKER AGAR BASE (IS: 5887 (Part II) 1976, reaffirmed 2005.) / TM 943 - BAIRD PARKER AGAR BASE W/ SULPHA along with 50 ml Concentrated Egg Yolk Emulsion or 10 ml in 1000 ml TM 118 - GIOLITTI-CANTONI BROTH BASE / TM 118A - GIOLITTI-CANTONI BROTH BASE (ISO 6888-3:2003) / TM 522 - HOYLE MEDIUM BASE along with 50 ml of laked blood. Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**

Revision: 11 March., 2022

