

## TS 024 – CLOSTRIDIUM DIFFICILE SUPPLEMENT

### INTENDED USE

For selective isolation and enumeration of *Vibrio* species from food.

### COMPOSITION

Ingredients	Concentration
D-Cycloserine	250mg
Cefoxitin	8mg

(Per vial sufficient for 500 ml medium)

### Applied in Medium

TM 068 - CLOSTRIDIUM DIFFICILE AGAR BASE













TM 2042 - CLOSTRIDIUM BRAZIER AGAR BASE

### INSTRUCTION FOR USE

Rehydrate the contents of one vial aseptically with 2 ml sterile distilled water. Mix well and aseptically add along with 7% v/v defibrinated horse blood to 500 ml sterile, molten, cooled (45-50°C) TM 068 - CLOSTRIDIUM DIFFICILE AGAR BASE / TM 2042 - CLOSTRIDIUM BRAZIER AGAR BASE. Mix well and pour into sterile petri plates. Sheep blood may be used in place of horse blood but some strains of the organism will show a slightly reduced growth.

### STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>Müller GmbH Bachstraße 16 41143 Aachen, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

**\*For Lab Use Only**

Revision: 8 March., 2022