

## TS 033 – FRASER ENRICHMENT SUPPLEMENT

### INTENDED USE

For isolation and enrichment of *Listeria monocytogenes* from food and environmental samples.

### COMPOSITION

Ingredients	Concentration
Acriflavine hydrochloride	25mg
Nalidixic acid	4mg

(per vial sufficient for 1000 ml medium)

### Applied in Medium










TM 113 - FRASER SECONDARY ENRICHMENT BROTH BASE

### INSTRUCTION FOR USE

Rehydrate the contents of one vial aseptically with 10 ml of sterile distilled water. Mix well and aseptically add it to 990 ml sterile, cooled (45-50°C) TM 113 - FRASER SECONDARY ENRICHMENT BROTH BASE. Mix well and dispense as desired.

### STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 Best Before	 QTY. Quantity	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 LOT/ B. NO. Lot / Batch Number	 Consults Instructions for Use	 QR Code	

**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

\*For Lab Use Only  
Revision: 28 Feb., 2022