

TS 043 – VANCLO T SUPPLEMENT (Vancomycin - Colistin - Amphotericin - B Trimethoprim)

INTENDED USE

For selective isolation of *Neisseria* species.

COMPOSITION

Ingredients	Concentration
Vancomycin	1mg
Colistin sulphate	3.750mg
Amphotericin B	0.500mg
Trimethoprim lactate	1.500mg

(per vial sufficient for 500 ml medium)

Applied in Medium

TM 116 - GC AGAR BASE













TMV 116 - GC AGAR BASE (VEG.)

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 10 ml sterile distilled water. Mix gently. Avoid frothing of the solution to dissolve. Aseptically add the contents of one vial to 490 ml sterile, molten, cooled (45-50°C) TM 116 - GC AGAR BASE / TMV 116 - GC AGAR BASE (VEG.) along with Haemoglobin Solution and GC Supplement w/ Antibiotics. Also add sterile rehydrated contents of 1 vial of Yeast Autolysate Supplement or Vitamin amino Growth Supplement. Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedWor GmbH Buckhorn 16, 41143 Krefeld, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 2 March., 2022