

## TS 057 – YERSINIA SELECTIVE SUPPLEMENT

### INTENDED USE

For selective isolation of *Yersinia enterocolitica*.

### COMPOSITION

Ingredients	Concentration
Cefsulodin	7.500mg
Triclosan(Irgasan)	2mg
Novobiocin	1.250mg

(per vial sufficient for 500 ml medium)

### Applied in Medium

TM 501 - YERSINIA SELECTIVE AGAR BASE (CIN AGAR)

TM 2439 - YERSINIA SELECTIVE AGAR BASE, W/ 1.2% AGAR













TM 2440 - YERSINIA SELECTIVE BROTH BASE

### INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 2 ml of sterile distilled water and 1 ml of ethanol. Mix gently to dissolve the contents completely and aseptically add to 500 ml of sterile, molten, cooled (45-50°C) TM 501 - YERSINIA SELECTIVE AGAR BASE (CIN AGAR) / TM 2439 - YERSINIA SELECTIVE AGAR BASE, W/ 1.2% AGAR / TM 2440 - YERSINIA SELECTIVE BROTH BASE. Mix well and pour into sterile petri plates.

### STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative MedWor GmbH Buckhorn 10, 40143 Essen, Germany	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

**\*For Lab Use Only**

Revision: 2 March., 2022