

TS 098 – WILLIS AND HOBBS SUPPLEMENT

INTENDED USE

For isolation of *Clostridium perfringens* and *Clostridium botulinum* species from food.

COMPOSITION

Ingredients	Concentration
Neomycin sulphate	33mg

(per vial sufficient for 475 ml medium)

Applied in Medium













TM 1319 - WILLIS AND HOBBS MEDIUM BASE (IS: 5887 (Part 4) : 1999)

INSTRUCTION FOR USE

Rehydrate the contents of one vial aseptically with 5.0 ml of sterile distilled water and aseptically add it to 400 ml of sterile, molten, cooled (45-50°C) TM 1319 - WILLIS AND HOBBS MEDIUM BASE (IS: 5887 (Part 4) : 1999) along with 15 ml of Egg Yolk Emulsion and 60 ml of sterile skim milk.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative	 CE European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 2 March., 2022