

TS 118 – LISTERIA UVM SUPPLEMENT-II

INTENDED USE

For selective isolation and cultivation of *Listeria monocytogenes* from clinical specimens.

COMPOSITION

Ingredients	Concentration
Acriflavin hydrochloride	12.500mg
Nalidixic acid	10mg

(Per vial sufficient for 500 ml medium)

Applied in Medium













TM 1225 - LISTERIA ENRICHMENT MEDIUM BASE (UVM)

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 2 ml of sterile distilled water. Mix well and aseptically add to 500 ml sterile, cooled (45-50°C) TM 1225 - LISTERIA ENRICHMENT MEDIUM BASE (UVM). Mix well and dispense as desired.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative MedWor GmbH Buckhorn 10, 40143 Düsseldorf, Germany	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 7 March., 2022