

TS 205 – CHROMOGENIC LISTERIA SELECTIVE SUPPLEMENT

INTENDED USE

For the isolation, detection and enumeration of *Listeria monocytogenes*.

COMPOSITION

Ingredients	Concentration
Ceftazidime	2mg
Amphotericin B	2.500mg

(Per vial sufficient for 500 ml medium)

Applied in Medium

TM 1845 - CHROMOGENIC L-MONO LISTERIA DIFFERENTIAL AGAR

TM 004 - CHROMOGENIC LISTERIA AGAR BASE (ISO 11290-1 & 2:2004, ISO 11133:2014)













TM 1634 - CHROMOGENIC LISTERIA AGAR BASE (Modified)

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial with 5 ml of sterile distilled water. Mix well and aseptically add to 500 ml sterile, molten, cooled (45-50°C) TM 1845 - CHROMOGENIC L-MONO LISTERIA DIFFERENTIAL AGAR / TM 004 - CHROMOGENIC LISTERIA AGAR BASE (ISO 11290-1 & 2:2004, ISO 11133:2014) / TM 1634 - CHROMOGENIC LISTERIA AGAR BASE (Modified). Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedMer GmbH Buckhorn 16, 41143 Alsdorf, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only
Revision: 8 March., 2022