

TS 248 – CHROMOGENIC EC O157:H7 SELECTIVE SUPPLEMENT

INTENDED USE

For selective isolation and detection of *Escherichia coli* O157:H7 from food samples.

COMPOSITION

Ingredients	Concentration
Novobiocin	10mg
Potassium tellurite	1mg

(Per vial sufficient for 1000 ml medium)

Applied in Medium

TM 1854 - CHROMOGENIC EC O157:H7 AGAR (ISO 16654:2001)

TM 2123 - CHROMOGENIC ENRICHMENT BROTH BASE FOR EC O157:H7













TM 2017 - BUFFERED PEPTONE WATER

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 10 ml of sterile distilled water. Mix well and add aseptically to 990 ml sterile, molten, cooled (45-50°C) TM 1854 - CHROMOGENIC EC O157:H7 AGAR (ISO 16654:2001) / TM 2123 - CHROMOGENIC ENRICHMENT BROTH BASE FOR EC O157:H7 / TM 2017 - BUFFERED PEPTONE WATER. Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative Mallat Gold Bachmann 10, 40143 Alsdorf, Germany	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 8 March., 2022