

TS 256 – STREPTOCOCCUS SELECTIVE SUPPLEMENT

INTENDED USE

Recommended for the selective isolation and cultivation of Streptococci of medical and veterinary importance.

COMPOSITION

Ingredients	Concentration
Colistin sulphate	5.000mg
Oxolinic acid	2.500mg

Applied in Medium

TM 2044 - COLUMBIA BLOOD AGAR BASE W/ 1% AGAR

TMV 071 - COLUMBIA BLOOD AGAR BASE (VEG.)













TM 071 - COLUMBIA BLOOD AGAR BASE

INSTRUCTION FOR USE

Rehydrate the contents of one vial aseptically with 1 ml of 0.2N NaOH and 4 ml of sterile distilled water. Mix well and aseptically add to 500 ml of sterile, molten, cooled (45-50°C) Columbia Blood Agar Base (TM 071) along with 5% v/v defibrinated blood. Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative MedNet GmbH Buckstrasse 10, 49163 Muenster, Germany	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 12 Oct., 2022